

Clathroporina Müll. Arg.
(TRICHOTHELIACEAE)

After various authors

Rev. 5/96

Thallus crustose, epi- or endo-phloic, uniform, ecorticate, amorphous, spreading in patches, closely attached to substrate at first but ultimately peeling or breaking from it, olive-green to yellowish brown. Perithecia immersed (\pm completely covered by thallus tissues; often in swollen pustules) to sessile (according to Rogers, but perhaps referring to the pustules), not aggregated; ostiole erect, straight, punctiform to \pm dilated, dark brown to black; exciple colorless or dark, hemispherical to \pm globose; paraphyses simple to sparingly branched, free; asci thin-walled, unitunicate, clavate, 1-, (2-)6-8 spored; spores ellipsoid to fusiform, muriform, hyaline, walls thin, of even thickness. Pycnidia immersed; fulcrum endobasidial; pycnospores oblong to oblong-cylindrical. Photobiont Trentepohlia. On bark or sometimes spreading onto shaded rocks, often in deeply shaded humid habitats, mainly tropical.

1. Thallus isidiate. Ascospores 8-celled, 32-48 x 5-6(-7.5) μ m.

Florida. C. isidiifera R. C. Harris

1. Thallus not isidiate. 2

2. Ascospores 8-celled, 40-55 x 7.5-9(-11) μ m. Florida. C. subpungens (Malme) R. C. Harris

2. Ascospores 8-celled, 35-50 x 5-7 μ m. Florida. C. tetracerae (Ach.) R. C. Harris

C. isidiifera R. C. Harris

Thallus light green fresh, grayer in herbarium, shiny, continuous, not separating from substrate, with a dark brown to black hypothallus, isidiate; isidia cylindrical to slightly clavate, occasionally branched, with a ca. 10-15 μ m cortical/algal layer and a ca. 25 μ m diam. core of oxalate crystals. Ascomata subglobose, 0.2-0.3 mm diam., immersed or in somewhat emergent warts; wart 0.4-0.7 mm diam.; wall colorless below to yellowish or orange-brown around ostiole. Asci cylindrical, ca. 140 x 12 μ m, with eight obliquely \pm biseriate spores. Ascospores cylindrical-fusiform, mostly \pm equally tapered at either end, 8-celled, 32-48 x 5-6(-7.5) μ m; microconidia not found.

C. mastoidea (Ach.) R. C. Harris (= misidentifications for N. America)

Thallus epilithic or epiphloedal, pale gray, pale khaki-gray to pale gray-green (white or whitish gray according to ?), color not changing when wet, or becoming dull greenish; continuous to areolate, sometimes peeling, matt to glossy, smooth to irregularly rugulose, (50-)100(-200) μ m thick, K+ dull

orange-brown or red-brown, heavily impregnated with crystal-like structures and/or minute rock fragments; areoles 0.2-0.8(-1.0) mm wide, irregular, angular or rounded, convex, frequently rimulose; upper layer usually colorless, 8-14 μm thick, of closely packed, 2-3(-4) μm thick, prosoplectenchymatous hyphae; algal layer with vertically or diagonally aligned 3-4 μm wide hyphae; basal layer 15-30 μm thick, colorless to brown-black, composed of cells similar in size and alignment to those of cortex. Prothallus whitish, brown-black or not apparent. Algae broadly ellipsoid to globose, 8-16 x 8-13 μm .

Perithecia immersed in thallus-dominated verrucae. Verrucae (0.36-)0.64(-0.9) mm diam., hemispherical, not or only slightly constricted at the base, numerous, usually solitary (or occasionally 2-3-aggregated); perithecial apex rounded or flattened; ostiole inconspicuous or in a 50-100 μm wide apical papilla or in a 40-80 μm wide depression, brown, blackening, K+ reddish; involucrellum not exposed, apical to dimidiate, contiguous with the excipulum (arching away from it according to ?), 0.15-0.20 mm diam., 20-60 μm thick; pale to medium brown except for a dull or glossy black, 0.05-0.2(-0.3) mm wide, 20-40 μm thick, area near the apex (outer layer dark yellowish brown, golden yellow within, according to ?), heavily impregnated with crystal-like structures and containing algal cells; excipulum (15-)20-25 μm thick, medium orange-brown towards the apex and at the sides, pale yellow-brown at the base; centrum globose to depressed-ovate, 0.25-0.44 mm diam. (0.15-0.20 mm according to ?), without oil globules, I-; paraphyses simple, 0.8-1.2 μm thick; periphyses absent; asci elongate-cylindrical, 8-spored, with rounded or truncate apices, 115-162 x 16-25 μm (100-110 x 6-9 μm according to ?); ascospores uniseriate or sometimes biseriate, Spores 7(-8)-septate, (32-)49(-65.5) x (6-)9(-12.5) μm [according to McCarthy; 5(-7)-septate, 32-50 x 5-7 μm , according to Awasthi; 7-15 μm wide according to Riddle; 19-30 x 4-7 μm according to ?], fusiform or elongate-cylindrical (or ellipsoid according to ?), straight, curved, or faintly sigmoid, irregularly biseriate in the asci; with pointed ends (according to ?); gelatinous sheath (1.5-)2-3(-3.5) μm thick (to 1 μm according to ?); contents clear.

Pycnidia usually numerous, semi-immersed to almost entirely immersed, (0.08-)0.12(-0.14) mm diam., brown-black above, colorless to pale brown below, with a richly convoluted conidogenous layer. Pycnospores fusiform to elongate-fusiform, 3-4(-5) x 1 μm .

On bark. Florida; in other parts of the world also reported from rock.

C. amygdalina = misidentification (N. American records) of Julella sublactea

C. diphloea = Laurera megasperma

C. exiguella = Julella sublactea (?--"almost certainly a Thelenella" according to McCarthy, 1995)

Literature

Awasthi. 19 . Microlichens of India, etc.

Fink, B. 1935. Lichen Flora of the United States.

Galloway, D. 1985. Flora of New Zealand Lichens.

Harris, R. C. 1995. More Florida Lichens.

McCarthy, P. M. 1995. A reappraisal of Clathroporina Müll. Arg. (Trichotheliaceae). Lichenologist 27(5): 321-350.

Rogers, 19_. Genera of Australian Lichens.

Tucker, S. and R. C. Harris. 1980.